arXiv:math/0505163v2 [math.DG] 27 May 2005

A note on uniformization of Riemann surfaces by Ricci flow

Xiuxiong Chen, Peng Lu and Gang Tian

In this note, we clarify that the Ricci flow can be used to give an independent proof of the uniformization theorem of Riemann surfaces. The key is the simple observation stated in Lemma 1.

When metric g_0 has positive curvature on S^2 or is a metric on a surface with genus bigger 1, Hamilton proves that the volume-normalized Ricci flow $g(t)$ with $g(0) = g_0$ converges to a metric of constant curvature [Ha], however the proof in §10 [Ha] of the fact that gradient shrinking soliton on S^2 has positive constant curvature, uses the uniformization theorem. There is another proof of the fact which uses the Kazdan-Warner identity (see p. 131 [CK]), but the proof of the Kazdan-Warner identity uses the uniformization theorem. When g_0 does not have positive curvature on S^2, Chow proves that $g(t)$ will have positive curvature for large t [Ch1], in the proof the entropy formula is proved using the uniformization theorem. Later different proofs of entropy estimate, which replaces the entropy formula, are given without using the uniformization theorem [Ch2], [CW]. So the only place, that the uniformization theorem is used for $g(t)$ to converge to a metric of constant curvature, is to argue that a gradient shrinking Ricci soliton on S^2 has positive constant curvature.

Lemma 1 *Let (Σ^2, g) be a two dimensional complete Riemannian manifold with non trivial Killing vector field X. Suppose X vanishes at $O \in \Sigma$, then (Σ^2, g) is rotationally symmetric.*

Proof. Let $\Phi_t : \Sigma \to \Sigma, t \in (-\infty, \infty)$ be the isometry group generating by X, $\frac{d}{dt}\Phi_t(x) = X(\Phi_t(x))$. By assumption $\Phi_t(O) = O$ for all t. Hence the tangent of Φ_t induces an oriented linear isometry

$$(\Phi_t)_* : (T_O\Sigma, g(O)) \to (T_O\Sigma, g(O)).$$

Since the oriented linear isometry group of $(T_O\Sigma, g(O))$ is S^1 and the map $t \to (\Phi_t)_*$ is a nontrivial homomorphism, there is a $t_0 > 0$ such that $(\Phi_0)_* = (\Phi_{t_0})_*$. Note that isometry Φ_t is determined by $(\Phi_t)_*$ through minimal geodesic starting at O, we have $\Phi_{t_0} = \Phi_0$. We have shown that there is a nontrivial S^1 isometric action on (Σ, g). It is clear that if there is a ray starting at O then Σ is topologically $\mathbb{R}^2$; If there is not any ray starting at O then Σ is topologically S^2. ■

It was known that a complete gradient steady gradient soliton in dimension two with positive curvature must be a cigar solution (see the Editors' footnote on p.241-242 [3CY]). Now using the rotational symmetry from Lemma 1, we can prove a similar proposition for gradient shrinking solitons without using the uniformization theorem.

Proposition 2 *If g is a gradient shrinking soliton on closed surface Σ^2, then g has positive constant curvature.*

Proof. Recall the gradient shrinking soliton equation $R_{ij} = cg_{ij} + \nabla_i \nabla_j f$ for $c > 0$ and some smooth function f on Σ. We choose $c = 1$ below. Since Σ is closed, there is a point O such that $\nabla f(O) = 0$. Since $R_{ij} = \frac{1}{2} R g_{ij}$ in dimension 2, it follows from soliton equation that ∇f is a conformal vector field. Let J be the almost complex structure on $T\Sigma$ defined by 90° counterclockwise rotation. It is well-known that $J(\nabla f)$ is a Killing vector field (see, for example, the Editor's footnote on p.241-242 [3CY]). Hence by Lemma 1 g is rotationally symmetric $g = dr^2 + h(r)^2 d\theta^2, 0 \le r \le A < \infty, 0 \le \theta \le 2\pi$. From the proof of Lemma 1 we may assume $f = f(r)$. Using Gauss curvature $K_g = -\frac{h''}{h}$ the soliton equation becomes

$$-\frac{h''}{h} = 1 + f'' \qquad -\frac{h''}{h} = 1 + \frac{h'f'}{h}.$$

Integrating $f'' = \frac{h'f'}{h}$ we get $f' = ah$ for some constant a. Hence $-\frac{h''}{h} = 1 + ah'$, multiplying hh' and integrating over $[0, A]$, we get

$$-\frac{(h')^2}{2}\Big|_0^A = \frac{h^2}{2}\Big|_0^A + a\int_0^A h(h')^2 dr.$$

It follows from the metric $dr^2 + h(r)^2 d\theta^2$ is smooth on Σ at $r = 0$ and $r = A$ that $h(0) = h(A) = 0$ and $h'(0) = -h'(A) = 1$. So $a = 0$ and g is Einstein metric. ■

Hamilton proved that any possible nontrivial soliton on closed surface must be a gradient shrinking soliton (see the proof in §10.1 [Ha]). So there are no nontrivial solitons on closed surfaces.

In higher dimensions, one may have nontrivial shrinking Ricci solitons. However, X. C and G. T can prove without using the uniformization theorem that any shrinking Kähler-Ricci soliton with positive bisectional curvature must be Kähler-Einstein [CT].

P.L thanks Bennett Chow for discussion about the Editor's note in [3CY]. Authors are supported by NSF research grants.

References

[3CY] H.D. Cao, B. Chow, S.C. Chu and S.T. Yau (editors), *Collected papers on Ricci flow*. Internat. Press, Somerville, M.A. (2003).

[Ch1] B. Chow, *The Ricci flow on the 2-sphere*. J. Diff. Geom. **33** (1991), 325-334.

[Ch2] B. Chow, *On the entropy estimate for the Ricci flow on compact 2-orbifolds*. J. Diff. Geom. **33** (1991), 597-600.

[CK] B. Chow and D. Knopf, *The Ricci flow: An introduction*. AMS, Providnece, RI, (2004).

[CT] X.X Chen and G. Tian, unpublished notes.

[CW] B. Chow and L.F. Wu, *The Ricci flow on compact 2-orbifolds with curvature negative somewhere*. Comm. Pure Appl. Math. **44** (1991), 275-286.

[Ha] R. Hamilton, *The Ricci flow on surfaces*, Contemp. Math. **71** AMS, Providnece, RI, (1988), 237-262.

Xiuxiong Chen, Dept of Math, University of Wisconsin, Madison, WI 53706

Peng Lu, Dept of Math, University of Oregon, Eugene, OR 97403

Gang Tian, Dept of Math, Princeton University, Princeton, NJ 08544